

Snapshot®



NURSERY



ORNAMENTAL

SPECIALTY HERBICIDE

Time is money. For nursery operators, landscapers and other turf and ornamental professionals, time spent on hand weeding is simply time taken from other important tasks.

Free up some of that valuable time with Snapshot® specialty herbicide. Snapshot delivers season-long control of more broadleaf and grassy weeds than any other preemergence herbicide. Snapshot works around the clock for up to six to eight months to keep landscapes and container- and field-grown nursery ornamentals weed-free. Snapshot has been tested on hundreds of ornamental plants and grasses and has been proven to offer exceptional plant tolerance.



How Snapshot® specialty herbicide Works

1 SNAPSHOT *Disrupts*
WEED SEED GERMINATION



1/2"

ACTIVATE WITH AT LEAST 1/2 INCH RAINFALL OR IRRIGATION

Snapshot bonds tightly with soil particles and is low in water solubility, so it won't move out of the application zone.

2 SEEDLINGS *Die Before*
BREAKING SOIL SURFACE



3 DAYS

RAINFALL OR IRRIGATION TO OCCUR WITHIN 3 DAYS FOR OPTIMAL RESULTS

Snapshot® specialty herbicide
controls or suppresses

125 BROADLEAF WEEDS
& ANNUAL GRASSES

including:

- | | |
|-------------|------------------|
| Spurge | Annual bluegrass |
| Dandelion | Crabgrass |
| Bittercress | Henbit |
| Chickweed | Marestail |
| Oxalis | |

Nursery Applications

*over-the-top application
on ornamentals*



Application Precautions

- Snapshot does not control established weeds. Apply prior to weed seed germination. Control existing weeds by cultivation or with a postemergence herbicide.
- Weed residues, prunings and debris should be removed or thoroughly mixed into soil prior to treatment.
- Repeat applications at the rate of 150 pounds per acre and higher should not be made within 60 days of original application. Do not apply more than 600 pounds per acre within a 12-month period.
- Do not apply Snapshot to:
 - Newly transplanted ornamentals (until soil has been settled by packing and irrigation or rainfall)
 - Un-rooted liners or cuttings that have been planted in pots for the first time
 - Pots less than 4 inches wide

When to Apply



EARLY SPRING PRIOR TO
GERMINATION OF TARGET WEEDS

LATE SUMMER TO EARLY FALL

ANY TIME IMMEDIATELY AFTER CULTIVATION

Following application, monitor weed control over time to determine if additional applications may be needed.

Active Ingredients

Snapshot® specialty herbicide contains the active ingredients isoxaben and trifluralin. Together, they provide excellent plant tolerance and long-lasting, broad-spectrum preemergence weed control.


Available Formulations

Snapshot is available in two easy-to-use granular formulations — Snapshot 2.5 TG and Snapshot DG specialty herbicides. Available in convenient 25-pound bags, Snapshot DG utilized The Andersons patented DG Lite formulation which features a dispersible granule.

Additional Information

Contact the Customer Information Center at Dow AgroSciences via email (info@dow.com), fax (800.905.7326) or phone (800.255.3726) for more information. Or contact your local Dow AgroSciences sales representative.

How to Apply



Depending on weeds to be controlled, apply at labeled rate:

2.3 - 4.6 POUNDS *per*
1,000 SQUARE FEET

DowProvesIt.com



Dow AgroSciences

Solutions for the Growing World

