

US79397887F (121114Fv2) LABMC BELT SC (DEU) 480 1 GALLON

| LEGUME VEGETABLES Except SOYBEAN | |
|---|---|
| <p>Crops of Crop Groups 6 and 7 including Edible-podded and Succulent Shelled Pea and Bean, Dried Shelled Pea and Bean and Foliage of Legume Vegetables:</p> <p>Bean (<i>Lupinus</i> spp., includes grain lupin, sweet lupin, white lupin, white sweet lupin)</p> <p>Bean (<i>Phaseolus</i> spp., includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean)</p> <p>Bean (<i>Vigna</i> spp., includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, Crowder pea, moth bean, mung bean, rice bean, Southern pea, Urd bean, yardlong bean)</p> <p>Pea (<i>Pisum</i> spp., includes dwarf pea, edible-pod pea, English pea, field pea, garden pea, green pea, snow pea, sugar snap pea)</p> <p>Other Peas and Beans: Broad bean (fava bean), chickpea (garbanzo bean), guar, jackbean, lablab bean (hyacinth bean), lentil, pigeon pea, sword bean</p> | |
| PESTS CONTROLLED | RATE PER APPLICATION fluid oz/Acre |
| <p>Alfalfa caterpillar Alfalfa looper Armyworm Beet armyworm Cabbage looper Celery looper Corn earworm Cutworm species European corn borer Fall armyworm Green cloverworm Imported cabbageworm Leaf skeletonizer species Leaf-tier species</p> | <p>Lesser cornstalk borer Painted lady (thistle) caterpillar Saltmarsh caterpillar Silverspotted skipper Southern armyworm Southwestern corn borer Soybean looper Tobacco budworm Velvetbean caterpillar Webworm species Western bean cutworm Wollybear caterpillar Yellowstriped armyworm Western yellowstriped armyworm</p> |
| <p>2.0 - 3.0</p> | |
| <p>Notes and Restrictions</p> <p>Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours.</p> <p>Pre-harvest Interval (PHI): Edible podded and succulent shelled peas and beans - 1 day; Dry peas and beans – 14 days; Forage, hay and vines – 3 days.</p> <p>Do not apply more than 3.0 fl oz per acre (0.094 lb ai/A) per 5 day interval.</p> <p>Do not apply more than 6.0 fl oz per acre (0.188 lb ai/A) per crop season.</p> <p>Apply BELT SC Insecticide in sufficient water volume that provides thorough coverage of plant foliage and fruit.</p> <p>Minimum application volume: 10.0 GPA – ground; 2.0 GPA – aerial application</p> <p>See CHEMIGATION statement in <i>Application Guidelines</i> section of this label.</p> | |

US79397887F (121114Fv2) LABMC BELT SC (DEU) 480 1 GALLON

| PEANUT | | RATE PER APPLICATION fluid oz/Acre |
|--|------------------------|---|
| PESTS CONTROLLED | | |
| Armyworm | Fall armyworm | 2.0 - 4.0 |
| Beet armyworm | Loopers | |
| Corn earworm | Rednecked peanutworm | |
| Cutworms | Southern armyworm | |
| Green cloverworm | Velvetbean caterpillar | |
| Notes and Use Restrictions | | |
| Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours. | | |
| Pre-harvest Interval (PHI): 3 days . | | |
| Do not apply more than 4.0 fl oz per acre (0.125 lb ai/A) per 7-day interval . | | |
| Do not apply more than 12.0 fl oz per acre (0.375 lb ai/A) per crop season . | | |
| Minimum application volume: 10.0 GPA – ground; 2.0 GPA – aerial application | | |
| See CHEMIGATION statement in <i>Application Guidelines</i> section of this label. | | |

| POME FRUIT | | RATE PER APPLICATION fluid oz/Acre | |
|---|-----------------------------|---|--|
| Crops of Crop Groups 11 including: Apple, Crabapple, Loquat, Mayhaw, Oriental pear, Pear, Quince | | | |
| PESTS CONTROLLED | | | |
| Codling moth (West of the Rockies) | | 5.0 | |
| <i>For use against low to moderate infestations in conjunction with alternate control measures such as in established mating disruption blocks.</i> | | | |
| Codling moth (East of the Rockies) | Oriental fruit moth | 3.0 - 5.0 | |
| Eyespotted bud moth | Pandemis leafroller | | |
| Fall webworm | Redbanded leafroller | | |
| Fruitree leafroller | Spotted tentiform leafminer | | |
| Green fruitworm | Tufted apple bud moth | | |
| Lacanobia fruitworm | Variigated leafroller | | |
| Lesser appleworm | Western tentiform leafminer | | |
| Obliquebanded leafroller | | | |
| Notes and Use Restrictions | | | |
| Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours. | | | |
| Pre-harvest Interval (PHI): 14 days . | | | |
| Do not apply more than 5.0 fl oz per acre (0.156 lb ai/A) per 7-day interval . | | | |
| Do not apply more than 15.0 fl oz per acre (0.468 lb ai/A) per crop season . | | | |
| Do not apply more than 3 times per crop season. | | | |
| Apply BELT SC Insecticide in sufficient water volume that provides thorough coverage of plant foliage and fruit. | | | |
| Aerial application is prohibited. | | | |

US79397887F (121114Fv2) LABMC BELT SC (DEU) 480 1 GALLON

| SOYBEAN | | | RATE PER APPLICATION fluid oz/Acre |
|--|---------------------------|------------------------|---|
| PESTS CONTROLLED | | | |
| Alfalfa caterpillar | Imported cabbageworm | Tobacco budworm | 2.0 - 3.0 |
| Armyworm | Leaf skeletonizer species | Tobacco hornworm | |
| Beet armyworm | Lesser cornstalk borer | Tomato hornworm | |
| Cabbage looper | Painted lady (thistle) | Velvetbean caterpillar | |
| Corn earworm | caterpillar | Webworm species | |
| Cutworm species | Saltmarsh caterpillar | Wollybear caterpillar | |
| European corn borer | Silverspotted skipper | Yellowstriped | |
| Fall armyworm | Southern armyworm | armyworm | |
| Green cloverworm | Soybean looper | | |
| Notes and Use Restrictions | | | |
| Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours. Pre-harvest Interval (PHI): Immature seed – 1 day ; Dry seed - 14 days ; Forage and hay – 3 days . Do not apply more than 3.0 fl oz per acre (0.094 lb ai/A) per 5-day interval . Do not apply more than 6.0 fl oz per acre (0.188 lb ai/A) per crop season . Minimum application volume: 10.0 GPA – ground; 2.0 GPA – aerial application See CHEMIGATION statement in <i>Application Guidelines</i> section of this label. | | | |

| SORGHUM | | | RATE PER APPLICATION fluid oz/Acre |
|--|-------------------------|--|---|
| PESTS CONTROLLED | | | |
| Armyworm | Sorghum webworm | | 2.0 - 4.0 |
| Beet armyworm | Southern armyworm | | |
| Cutworms | Southwestern corn borer | | |
| European corn borer | Stalk borer | | |
| Fall armyworm | Sugarcane borer | | |
| Mexican rice borer | Webworms | | |
| Sorghum headworm | Yellowstriped armyworm | | |
| Notes and Use Restrictions | | | |
| Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours. Pre-harvest Interval (PHI): Forage – 3 days ; grain and stover – 14 days . Do not apply more than 4.0 fl oz per acre (0.125 lb ai/A) per 7-day interval . Do not apply more than 12.0 fl oz per acre (0.375 lb ai/A) per crop season . Minimum application volume: 10.0 GPA – ground; 2.0 GPA – aerial application See CHEMIGATION statement in <i>Application Guidelines</i> section of this label. | | | |

US79397887F (121114Fv2) LABMC BELT SC (DEU) 480 1 GALLON

| STRAWBERRY and LOW GROWING BERRY SUBGROUP (except cranberry) | |
|---|---|
| Crops of Crop Subgroup 13-07G (except cranberry) including: Bearberry, Bilberry, Blueberry (lowbush), Cloudberry, Lingonberry, Muntries, Partridgeberry, Strawberry, plus cultivars, varieties and/or hybrids of these | |
| PESTS CONTROLLED | RATE PER APPLICATION fluid oz/Acre |
| Armyworm Corn earworm Cutworm Lesser cornstalk borer Omnivorous leaftier Strawberry leafroller | 2.0 – 2.4 |
| Notes and Use Restrictions Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours. Pre-harvest Interval (PHI): 8 days. Do not apply more than 2.4 fl oz per acre (0.075 lb ai/A) per 3-day interval. Do not apply more than 7.2 fl oz per acre (0.225 lb ai/A) per crop season. Minimum application volume: 10.0 GPA - ground, 2.0 GPA - aerial application. See CHEMIGATION statement in <i>Application Guidelines</i> section of this label. | |

US79397887F (121114Fv2) LABMC BELT SC (DEU) 480 1 GALLON

| STONE FRUIT | |
|--|--|
| Crops of Crop Group 12 including: Apricot, Cherry [sweet and tart], Nectarine, Peach, Plum [includes Chickasaw plum, Damson plum, and Japanese plum], Plumcot, Prune (fresh) | |
| PESTS CONTROLLED | RATE PER APPLICATION fluid oz/Acre |
| Codling moth Cherry fruitworm Eyespotted bud moth Fruittree leafroller Green fruitworm Lesser appleworm Obliquebanded leafroller Omnivorous leafroller Oriental fruit moth | Pandemis leafroller Peach twig borer Redbanded leafroller Redhumped caterpillar Spotted tentiform leafminer Threelined leafroller Tufted apple bud moth Variegated leafroller |
| 3.0 - 4.0 | |
| Notes and Use Restrictions Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours. Pre-harvest Interval (PHI): 7 days . Do not apply more than 4.0 fl oz per acre (0.125 lb ai/A) per 7-day interval . Do not apply more than 12.0 fl oz per acre (0.375 lb ai/A) per crop season . Do not apply more than 3 times per crop season. Apply BELT SC Insecticide in sufficient water volume that provides thorough coverage of plant foliage and fruit. Aerial application is prohibited. | |

| SUGARCANE | |
|---|---|
| PESTS CONTROLLED | RATE PER APPLICATION fluid oz/Acre |
| Sugarcane borer Mexican rice borer | |
| 3.0 – 4.0 | |
| Notes and Use Restrictions Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours. Pre-harvest Interval (PHI): 14 days . Do not apply more than 4.0 fl oz per acre (0.125 lb ai/A) per 7-day interval . Do not apply more than 12.0 fl oz per acre (0.375 lb ai/A) per crop season . Minimum application volume: 10.0 GPA – ground; 2.0 GPA – aerial application See CHEMIGATION statement in <i>Application Guidelines</i> section of this label. | |

US79397887F (121114Fv2) LABMC BELT SC (DEU) 480 1 GALLON

| SUNFLOWER and SAFFLOWER | |
|---|---|
| PESTS CONTROLLED | RATE PER APPLICATION fluid oz/Acre |
| Banded sunflower moth Cutworms Sunflower bud moth Sunflower moth Thistle caterpillar | 2.0 - 4.0 |
| <p>Notes and Use Restrictions Do not allow grazing or feed forage to livestock. Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours. Pre-harvest Interval (PHI): 14 days. Do not apply more than 4.0 fl oz per acre (0.125 lb ai/A) per 7-day interval. Do not apply more than 12.0 fl oz per acre (0.375 lb ai/A) per crop season. Minimum application volume: 10.0 GPA – ground; 2.0 GPA – aerial application</p> | |

| TOBACCO | |
|--|--|
| PESTS CONTROLLED | RATE PER APPLICATION fluid oz/Acre |
| Armyworm Beet armyworm Cabbage looper Corn earworm Cutworm species Fall armyworm Saltmarsh caterpillar | Southern armyworm Tobacco budworm Tobacco hornworm Tobacco splitworm Tomato hornworm Webworm species Yellowstriped armyworm 2.0 - 3.0 |
| <p>Notes and Use Restrictions Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours. Pre-harvest Interval (PHI): 14 days. Do not apply more than 3.0 fl oz per acre (0.094 lb ai/A) per 5-day interval. Do not apply more than 12.0 fl oz per acre (0.375 lb ai/A) per crop season. Do not apply more than 4 times per crop season. Minimum application volume: 10.0 GPA – ground; 2.0 GPA – aerial application See CHEMIGATION statement in <i>Application Guidelines</i> section of this label.</p> | |

US79397887F (121114Fv2) LABMC BELT SC (DEU) 480 1 GALLON

| TREE NUT CROPS | | | |
|---|--------------------------|-----------------------|---|
| Crops of Crop Group 14 and Pistachio including: Almond, Beech Nut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (hazelnut), Hickory Nut, Macadamia Nut, Pecan, Pistachio, Walnut (black and English) | | | |
| PESTS CONTROLLED | | | RATE PER APPLICATION fluid oz/Acre |
| Codling moth | Hickory shuckworm | Peach twig borer | 3.0 - 4.0 |
| Fall webworm | Naval orangeworm | Pecan nut casebearer | |
| Filbertworm | Obliquebanded leafroller | Redhumped caterpillar | |
| Fruittree leafroller | Omnivorous leafroller | Walnut caterpillar | |
| | | | |
| Notes and Use Restrictions | | | |
| Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours. | | | |
| Pre-harvest Interval (PHI): 14 days. | | | |
| Do not apply more than 4.0 fl oz per acre (0.125 lb ai/A) per 7-day interval. | | | |
| Do not apply more than 12.0 fl oz per acre (0.375 lb ai/A) per crop season. | | | |
| Apply BELT SC Insecticide in sufficient water volume that provides thorough coverage of plant foliage and fruit. | | | |
| Aerial application is prohibited. | | | |

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE

Do not store for more than 30 consecutive days at an average daily temperature exceeding 100° F. If allowed to freeze, shake well to ensure the product is homogenous before use. Store in original container and out of the reach of children, preferable in a locked storage area. Avoid cross contamination with other pesticides.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times, then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

US79397887F (121114Fv2) LABMC BELT SC (DEU) 480 1 GALLON

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

Bayer

15/16"

4-1/8"

7/16"

NET CONTENTS: 1 GALLON

BELT® SC Insecticide

GROUP 28 INSECTICIDE

ACTIVE INGREDIENT: Flubendiamide (*N*²-[1,1-dimethyl-2-(methylsulfonyl)ethyl]-3-iodo-*N*¹-[2-methyl-4-[1,2,2,2-tetrafluoro-1-(trifluoromethyl)ethyl]phenyl]-1,2-benzenedicarboxamide) 39%

OTHER INGREDIENTS: 61%

BELT SC Insecticide contains 4 pounds **TOTAL:** 100% of flubendiamide per US gallon (480 grams per liter).

EPA Reg. No. 264-1025

STOP – Read the label before use
KEEP OUT OF REACH OF CHILDREN
CAUTION

FOR ADDITIONAL PRECAUTIONARY STATEMENTS:
 See Attached Booklet.

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY**
 Call 24 Hours A Day 1-800-334-7577
 For **PRODUCT USE** Information Call 1-866-99BAYER
 (1-866-992-2937)

FIRST AID

| | |
|--------------------------------|--|
| IF ON SKIN OR CLOTHING: | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice. |
| IF SWALLOWED: | <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Have person sip a glass of water if able to swallow. • Do not give anything by mouth to an unconscious person. |

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For medical emergencies, health concerns, or pesticide incidents, you may call the Bayer CropScience Emergency Response toll free number 24 hours a day at 1-800-334-7577.

NOTE TO PHYSICIAN: No specific antidote is known. Treat symptomatically.

PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

FOR ADDITIONAL PRECAUTIONARY STATEMENTS:
See attached label booklet on: Personal Protective Equipment (PPE), User Safety Recommendations, and Environmental Hazards

DIRECTIONS FOR USE: See attached booklet.
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
 Read entire label before using this product.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE

Do not store for more than 30 consecutive days at an average daily temperature exceeding 100° F. If allowed to freeze, shake well to ensure the product is homogenous before use. Store in original container and out of the reach of children, preferable in a locked storage area. Avoid cross contamination with other pesticides.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times, then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Bayer CropScience LP
 P.O. Box 12014, 2 T.W. Alexander Drive
 Research Triangle Park, North Carolina 27709
 BELT is a registered trademark of Bayer.
 ©2013 Bayer CropScience US79397887F 121114Fv2 02/13

4-3/4"



(01) 0 0785740 13459 7

5"

5.5"